

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

Comparison of Gene Expression in Taxane Sensitive vs. Resistant Tumors




FIGURE 1

Extreme Drug Resistance (EDR[®]) Assay




FIGURE 2

Oncogene Expression Based Cell Sorting




FIGURE 3

Cell Sorting of Tumor Derived Vascular Endothelial Cells





FIGURE 4

Annexin V Apoptosis Assay



Schematic Representation of the Annexin V Assay. Early in apoptosis, phosphatidylserine (PS) is translocated from the inner to outer surface of the plasma membrane. In the presence of Ca^{2+} , Annexin V has a high affinity for PS. In unfixed cells, Annexin V binds only to PS that has been exposed to the external milieu.

FIGURE 5

Taxane Sensitive and Resistance Subsets by Annexin V and Propidium Iodide Based Cell Sorting




FIGURE 6

Comparison of Induced Gene Expression by Taxane Sensitive vs Resistant Breast Cancer




FIGURE 7

Venn Diagram Isolation of Genes Related to In Vitro Taxane Response Class




FIGURE 8